

## Prostaglandins and Sex Steroids from Reptilian (*Podarcis sicula sicula*) Ovarian Follicles at Different Developmental Stages

ANNA GOBBETTI, MASSIMO ZERANI, MARIA MADDALENA DI FIORE<sup>1</sup>  
and VIRGILIO BOTTE<sup>1</sup>

Department of Molecular, Cellular, and Animal Biology, University  
of Camerino, 62032 Camerino MC, Italy, and <sup>1</sup>Department of  
Zoology, University of Naples, 80134 Naples, Italy

**ABSTRACT**—Prostaglandin F<sub>2α</sub> (PGF<sub>2α</sub>), prostaglandin E<sub>2</sub> (PGE<sub>2</sub>), progesterone, androgens, and estradiol-17β *in vitro* basal release by follicles of the oviparous lizard, *Podarcis s. sicula* was studied; in addition, the *in vitro* effect of PGF<sub>2α</sub> and PGE<sub>2</sub> on sex steroid release was evaluated. Follicles were divided according to the different vitellogenic developmental stages: pre-vitellogenic, early-vitellogenic, mid-vitellogenic and fully-grown. PGF<sub>2α</sub> and progesterone basal release was highest in fully-grown follicles; PGE<sub>2</sub> and estradiol basal release was highest in early-vitellogenic follicles; androgens basal release was detectable in mid-vitellogenic and fully-grown vitellogenic follicles only. PGF<sub>2α</sub> increased progesterone release by fully-grown follicles; PGE<sub>2</sub> increased estradiol release by all follicle types, except by early-vitellogenic ones. The present data suggest that PGF<sub>2α</sub> and PGE<sub>2</sub> exert different roles on follicles: PGF<sub>2α</sub> seems to induce ovulation through the mediation of progesterone, while PGE<sub>2</sub> seems to be implied in the start and the sustaining of oocyte vitellogenic development through the mediation of estradiol.

### INTRODUCTION

In mammals, prostaglandins (PGs) of both F and E series modulate the ovary function, including steroidogenesis [6, 9, 18, 29]. PGs intervene in ovulation and oviduct relax in birds [1, 30]. As regards oviparous reptiles, oviposition is correlated with a dramatic elevation of prostaglandin F (PGF) and prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) in the loggerhead turtle, *Caretta caretta* [20] and the tuatara, *Sphenodon punctatus* [19]; PGF<sub>2α</sub> exhibits luteolytic effects in the lizard, *Anolis carolinensis* [21], and the snapping turtle, *Chelydra serpentina* [24], determining a decline in plasma progesterone. Little is known about the involvement of PGs in the vitellogenic development of follicles in reptiles, and, in particular, nothing is known about the role of PGs on the reproductive processes of the oviparous lizard, *Podarcis sicula sicula*, even if the

reproductive cycle of this lizard has been well studied [2, 3, 10].

In this study we have compared the PGF<sub>2α</sub>, PGE<sub>2</sub>, progesterone, androgens, and estradiol-17β *in vitro* basal releases by follicles, at different vitellogenic developmental stages, of the lizard, *Podarcis s. sicula*; in addition, the *in vitro* effect of PGF<sub>2α</sub> and PGE<sub>2</sub> on progesterone, androgens, and estradiol-17β releases by the same follicles has been examined.

### MATERIALS AND METHODS

**Animals** The reproductive cycle of the female lizard, *Podarcis s. sicula*, living in the surroundings of Naples (Campania, Italy, 25–75 m above sea level) is here briefly described. From the end of summer to the beginning of the next spring (stasis), gonads and genital apparatus are quiescent; then, soon after emerging, the ovary and oviduct undergo recovery and reach their full growth from March to mid-May (recovery period);

ovulation and egg deposition occur from mid-May to July (ovulatory period); after deposition, gonads and oviduct regress (postovulatory period) [2, 3, 10].

Adult female lizards, *Podarcis s. sicula*, of this population were captured in May and transferred to laboratory terraria, maintained in ambient photothermal conditions (L/D: 15/9; 18°C), and fed with meal worms and fresh vegetables *ad libitum*.

***In vitro* incubations of follicles** The lizards were sacrificed by decapitation, the developmental stage of the reproductive organs was observed and the ovaries rapidly removed. Follicles were freed under dissection microscope and placed in cold Dulbecco's modified Eagle medium (DME, Sigma, USA) containing 10 mM Hepes, 1 mg Penicillin/ml and 2 mg Streptomycin/ml. Follicles were divided according to the different vitellogenic developmental stages: pre-vitellogenic (500–1000  $\mu\text{m}$   $\phi$ ), early-vitellogenic (1500–3000  $\mu\text{m}$   $\phi$ ), mid-vitellogenic (4000–6000  $\mu\text{m}$   $\phi$ ) and fully-grown (>8000  $\mu\text{m}$   $\phi$ ). The follicles were placed in multiple tissue culture plates (Beckton Dickinson, USA); each well contained one follicle. Six follicles of each type were divided into 3 experimental groups (each consisting of 2 wells). The experimental groups were: a) follicle with DME alone; b) follicle with DME plus PGF<sub>2 $\alpha$</sub>  (150 ng); c) follicle with DME plus PGE<sub>2</sub> (150 ng). The final volume of each well was 1 ml. Culture plates were wrapped with aluminium foil and incubated in a shaking water bath (32°C), set at 30 rpm. One well of each experimental group was removed, respectively, after 3 and 6 hr of incubation. The incubation medium samples were immediately stored at –20°C for later hormone determination. In addition, the experiment was repeated without follicles. Tests on 5 parallel incubation sets were carried out. Preliminary evidence led our choosing to the incubation conditions and the PGF<sub>2 $\alpha$</sub>  or PGE<sub>2</sub> minimum effective dose utilized in the present *in vitro* study (data not shown).

***PGF<sub>2 $\alpha$</sub> , PGE<sub>2</sub>, progesterone, androgens, and estradiol-17 $\beta$  determination*** PGF<sub>2 $\alpha$</sub>  and PGE<sub>2</sub> were determined in incubation medium samples by

radioimmunoassay (RIA) [12, 13]. The determinations were carried out on duplicate incubation medium samples (500  $\mu\text{l}$ ) that were extracted with 5 ml of diethyl ether for 4 min. The organic fractions were transferred into glass tubes and evaporated to dryness under a nitrogen stream. The extracts were resuspended with 1 ml of assay buffer and assayed. The recovery of labelled PGF<sub>2 $\alpha$</sub>  and PGE<sub>2</sub> were  $83.3 \pm 1.3\%$  and  $88.8 \pm 1.7\%$ , respectively. Parallelism among the standard curve in buffer, a standard curve in incubation medium (then extracted) and a serial dilution of a single medium sample (extracted) were constant.

Concentrations of progesterone, androgens and estradiol-17 $\beta$  in incubation media were determined by RIA in accordance with methods previously reported [7, 13].

The following sensitivities were recorded: PGF<sub>2 $\alpha$</sub>  13 pg (intraassay variability: 7.5%; interassay variability: 14.5%); PGE<sub>2</sub>, 15.5 pg (intraassay variability: 7.0%; interassay variability: 12.5%); progesterone, 9 pg (intraassay variability: 8.0%; interassay variability: 10.0%); androgens, 7.5 pg (intraassay variability: 6.0%; interassay variability: 11.5%); estradiol-17 $\beta$ , 7 pg (intraassay variability: 8.5%; interassay variability: 13.0%). PGF<sub>2 $\alpha$</sub> , progesterone, testosterone, and estradiol-17 $\beta$  antisera were provided by Dr. G. F. Bolelli and Dr. F. Franceschetti (CNR-Physiopathology of Reproduction Service, University of Bologna, Italy). The PGE<sub>2</sub> antiserum was purchased from Cayman Chemical (USA). Testosterone was not separated from 5 $\alpha$ -dihydrotestosterone; therefore, the antiserum used is not specific and the data are expressed as androgens. Tritiated PGF<sub>2 $\alpha$</sub> , PGE<sub>2</sub>, progesterone, testosterone, and estradiol-17 $\beta$  were purchased from Amersham International (UK) and non-radioactive PGF<sub>2 $\alpha$</sub> , PGE<sub>2</sub>, progesterone, testosterone, and estradiol-17 $\beta$  from Sigma.

***Statistics*** Data relative to each hormone were submitted to analysis of variance (ANOVA) followed by Duncan's multiple range test [8, 31]. Correlation coefficients followed the procedures of Scossiroli and Palenzona [28].

## RESULTS

PGF<sub>2α</sub> basal release was higher ( $P < 0.01$ ) in fully-grown follicles with respect to the other follicle types; PGF<sub>2α</sub> was lower ( $P < 0.01$ ) in pre-vitellogenic with respect to early-vitellogenic and mid-vitellogenic follicles (Fig. 1). PGE<sub>2</sub> basal release was higher ( $P < 0.01$ ) in early-vitellogenic with respect to the other follicle types; PGE<sub>2</sub> was

lower in pre-vitellogenic with respect to mid-vitellogenic and fully-grown follicles (Fig. 2). Progesterone basal release was higher ( $P < 0.01$ ) in fully-grown with respect to the other follicle types; progesterone was higher ( $P < 0.01$ ) in mid-vitellogenic with respect to the pre-vitellogenic and early-vitellogenic follicles; progesterone was higher ( $P < 0.01$ ) in early-vitellogenic with respect to pre-vitellogenic follicles (Fig. 3, upper panel).




FIG. 1. PGF<sub>2α</sub> basal release by follicles, at different vitellogenic developmental stages (see Materials and Methods), of the oviparous lizard, *Podarcis s. sicula*. Follicles: pre-vitellogenic (black bars), early-vitellogenic (white bars), mid-vitellogenic (grey bars), fully-grown (hatched bars). Each mean refers to 5 values  $\pm$  SD. a,  $P < 0.01$  vs pre-vitellogenic, early-vitellogenic and mid-vitellogenic follicles; b,  $P < 0.01$  vs pre-vitellogenic follicles (Duncan's multiple range test).




FIG. 2. PGE<sub>2</sub> basal release by follicles, at different vitellogenic developmental stages (see Materials and Methods), of the oviparous lizard, *Podarcis s. sicula*. Follicles: pre-vitellogenic (black bars), early-vitellogenic (white bars), mid-vitellogenic (grey bars), fully-grown (hatched bars). Each mean refers to 5 values  $\pm$  SD. a,  $P < 0.01$  vs pre-vitellogenic, mid-vitellogenic and fully-grown follicles; b,  $P < 0.01$  vs pre-vitellogenic follicles (Duncan's multiple range test).




FIG. 3. Progesterone basal release (upper panel) and PGF<sub>2α</sub> effects (lower panel) on progesterone release by follicles, at different vitellogenic developmental stages (see Materials and Methods), of the oviparous lizard, *Podarcis s. sicula*. Follicles: pre-vitellogenic (black bars), early-vitellogenic (white bars), mid-vitellogenic (grey bars), fully-grown (hatched bars). Each mean refers to 5 values  $\pm$  SD. a,  $P < 0.01$  vs pre-vitellogenic, early-vitellogenic and mid-vitellogenic follicles; b,  $P < 0.01$  vs pre-vitellogenic and early-vitellogenic follicles; c,  $P < 0.01$  vs pre-vitellogenic follicles; \* $P < 0.01$  vs the same follicle type incubated with PGF<sub>2α</sub> (Duncan's multiple range test).

Androgens basal release was not detectable in pre-vitellogenic and early-vitellogenic follicles; androgens were higher ( $P < 0.01$ ) in fully-grown with respect to mid-vitellogenic follicles (Fig. 4). Estradiol-17 $\beta$  basal release was higher ( $P < 0.01$ ) in

early-vitellogenic with respect to the other follicle types; estradiol was lower ( $P < 0.01$ ) in pre-vitellogenic with respect to mid-vitellogenic and fully-grown follicles (Fig. 5, upper panel). PGF<sub>2α</sub> basal release values were positively correlated

FIG. 5. Estradiol-17 $\beta$  basal release (upper panel) and PGE<sub>2</sub> effects (lower panel) on estradiol-17 $\beta$  release by follicles, at different vitellogenic developmental stages (see Materials and Methods), of the oviparous lizard, *Podarcis s. sicula*. Follicles: pre-vitellogenic (black bars), early-vitellogenic (white bars), mid-vitellogenic (grey bars), fully-grown (hatched bars). Each mean refers to 5 values  $\pm$  SD. a,  $P < 0.01$  vs pre-vitellogenic, mid-vitellogenic and fully-grown follicles; b,  $P < 0.01$  vs pre-vitellogenic follicles; \* $P < 0.01$  vs the same follicle type incubated with PGE<sub>2</sub> (Duncan's multiple range test).






FIG. 4. Androgens basal release by follicles, at different vitellogenic developmental stages (see Materials and Methods), of the oviparous lizard, *Podarcis s. sicula*. Follicles: pre-vitellogenic (black bars), early-vitellogenic (white bars), mid-vitellogenic (grey bars), fully-grown (hatched bars). Each mean refers to 5 values  $\pm$  SD. a,  $P < 0.01$  vs mid-vitellogenic follicles (Duncan's multiple range test); nd, not detectable.





TABLE 1. Correlation coefficients among  $\text{PGF}_{2\alpha}$ ,  $\text{PGE}_2$ , progesterone and estradiol- $17\beta$  levels released by follicles, at different vitellogenic developmental stages, of *Podarcis s. sicula*

	Incubation times	
	3 hr	6 hr
$\text{PGF}_{2\alpha}$ vs progesterone	0.804	0.814
$\text{PGE}_2$ vs estradiol- $17\beta$	0.862	0.911

All correlations show the same level of significance ( $P < 0.001$ );  $\text{df} = 18$ .

( $P < 0.001$ ) to those of progesterone (Table 1);  $\text{PGE}_2$  basal release values were positively correlated ( $P < 0.001$ ) to those of estradiol- $17\beta$  (Table 1).

$\text{PGF}_{2\alpha}$  induced an increase ( $P < 0.01$ ) in progesterone release by fully-grown follicles (Fig. 3, lower panel).  $\text{PGE}_2$  induced an increase ( $P < 0.01$ ) in estradiol release by pre-vitellogenic, mid-vitellogenic, and fully-grown follicles (Fig. 5, lower panel).

## DISCUSSION

This is the first study regarding the *in vitro* basal release of PGs and sex steroids by the follicles, kept at different vitellogenic developmental stages, of the oviparous lizard *Podarcis s. sicula*.

$\text{PGF}_{2\alpha}$  showed the highest amounts in fully-grown follicle incubation samples; this finding supports the idea that  $\text{PGF}_{2\alpha}$  could be involved in the stimulation of ovulation; on the other hand, Goetz [15] reported that in several fish species, PGs can induce ovulation; in particular, plasma and ovarian tissue  $\text{PGF}_{2\alpha}$  levels are higher during ovulation in the yellow perch, *Perca flavescens* [17], in the brook trout, *Salvelinus fontinalis* [16], in the goldfish, *Carassius auratus* [4], and in the pond loach, *Misgurnus anguillicaudatus* [25]. Also in amphibians,  $\text{PGF}_{2\alpha}$  seems to be implied in ovulation processes, seeing that  $\text{PGF}_{2\alpha}$  *in vitro* induces ovulation in *Rana pipiens* [27]; more recently Gobbetti and Zerani [14] suggested that  $\text{PGF}_{2\alpha}$  is implied in the control of ovulation in *Rana esculenta*. The involvement of  $\text{PGF}_{2\alpha}$  in the ovulatory process of this lizard is also supported by the stimulatory role exerted by this prostaglandin on

progesterone increase by fully-grown follicles. Progesterone basal release was higher in fully-grown follicles, if compared with the other follicle types, so suggesting that this steroid is involved in the ovulation of *Podarcis*, as many authors reported for amphibian species [23]; in fact, progesterone, produced by follicle cells, induces ovulation in *Xenopus laevis* [32] and *Rana pipiens* [26].

The early-vitellogenic follicles released the highest values of  $\text{PGE}_2$  and estradiol, which both, however, remained still high in mid-vitellogenic and fully-grown follicles. The positive correlation between the basal release values of  $\text{PGE}_2$  and estradiol suggests a causal relationship and this hypothesis is validated by the  $\text{PGE}_2$  effect in increasing estradiol release. This finding might indicate a possible trigger role for this prostaglandin in the start of the vitellogenic development of the oocytes and also a role in sustaining the following vitellogenic growth of oocytes. In this context we recall that estradiol induces the hepatic vitellogenin synthesis for the development of the follicle, as widely reported for reptiles [5, 22]. The  $\text{PGE}_2$  stimulatory effect on estradiol release was detected in all follicle types, except for early-vitellogenic ones. This phenomenon could be due to the refractoriness of this follicle type to an additional stimulation of  $\text{PGE}_2$ ; in fact the early-vitellogenetic follicles released the highest amounts of  $\text{PGE}_2$  and estradiol.

Androgens were detected only in the incubation media of mid-vitellogenic and fully-grown follicles in accordance with Fortune [11], who found that only large follicles released these hormones *in vitro* in the amphibian *Xenopus laevis*. The meaning of this last result is unclear; it could be due to the catabolism of progesterone which exhibits the highest values in incubation media of mid-vitellogenic and fully-grown follicles; however, a role for androgens produced by these two follicle types cannot be excluded.

Summarizing, this work suggests different roles for  $\text{PGF}_{2\alpha}$  and  $\text{PGE}_2$  released by *Podarcis s. sicula* follicles:  $\text{PGF}_{2\alpha}$  could induce ovulation through progesterone mediation, while  $\text{PGE}_2$  could be implied in the start and the sustaining of vitellogenic development of oocytes through estradiol mediation.

## ACKNOWLEDGMENTS

The authors would like to thank James Burge, of the Camerino University Institute of Linguistics, for help with English. This work was supported by grant from MURST.

## REFERENCES

- Asem EK, Todd H, Hertelendy F (1987) *In vitro* effect of prostaglandins on the accumulation of cyclic AMP in the avian oviduct. *Gen Comp Endocrinol* 66: 244–247
- Botte V, Angelini F, Picariello O, Molino R (1976) The regulation of the reproductive cycle of the female lizard, *Lacerta sicula sicula* Raf. *Monitore Zool Ital* 10: 119–133
- Botte V, Delrio G (1965) Ricerche istochimiche sulla distribuzione dei 3- e 17-chetosteroidi e di alcuni enzimi della steroidogenesi nell' ovario di *Lacerta sicula*. *Boll Zool* 32: 191–198
- Bouffard RE (1979) The role of prostaglandins during sexual maturation, ovulation and spermiation in the goldfish, *Carassius auratus*. M Sc Thesis, University of British Columbia, Vancouver, BC, Canada
- Carnevali O, Mosconi G, Angelini F, Limatola E, Ciarcia G, Polzonetti-Magni A (1991) Plasma vitellogenin and 17 $\beta$ -estradiol levels during the annual reproductive cycle of *Podarcis s. sicula* Raf. *Gen Comp Endocrinol* 84: 337–343
- Dharmarajan AM, Sueoka K, Miyazaki S, Atlas SJ, Ghodgaonkar RB, Dubin NH, Zirkin BR, Wallach EE (1989) Prostaglandins and progesterone secretion in the *in vitro* perfused pseudopregnant rabbit ovary. *Endocrinology* 124: 1198–1203
- D'Istria M, Delrio G, Botte V, Chieffi G (1974) Radioimmunoassay of testosterone, 17 $\beta$ -oestradiol and estrone in the male and female plasma of *Rana esculenta* during sexual cycle. *Ster Lip Res* 5: 42–48
- Duncan DB (1955) Multiple range and multiple F test. *Biometrics* 11: 1–42
- Espey LL, Norris C, Saphire D (1986) Effect of time and dose of indomethacin on follicular prostaglandins and ovulation in the rabbit. *Endocrinology* 119: 746–754
- Filosa S (1973) Biological and cytological aspects of the ovarian cycle in *Lacerta sicula* Raf. *Monitore Zool Ital* 7: 151–165
- Fortune JE (1983) Steroid production by *Xenopus* ovarian follicles at different developmental stages. *Dev Biol* 99: 502–509
- Gobbetti A, Zerani M (1991) Gonadotropin-releasing hormone stimulates biosynthesis of prostaglandin F<sub>2 $\alpha$</sub>  by the interrenal gland of the water frog, *Rana esculenta*, *in vitro*. *Gen Comp Endocrinol* 84: 434–439
- Gobbetti A, Zerani M (1992) PGF<sub>2 $\alpha$</sub> , PGE<sub>2</sub> and sex steroids from the abdominal gland of the male crested newt *Triturus cristatus* (Laur.). *Prostaglandins* 43: 101–109
- Gobbetti A, Zerani M (1992) A possible involvement of prostaglandin F<sub>2 $\alpha$</sub>  (PGF<sub>2 $\alpha$</sub> ) in *Rana esculenta* ovulation: effects of mammalian gonadotropin-releasing hormone on *in vitro* PGF<sub>2 $\alpha$</sub>  and 17 $\beta$ -estradiol production from ovary and oviduct. *Gen Comp Endocrinol* 87: 163–170
- Goetz FW (1983) Hormonal control of oocyte final maturation and ovulation in fishes. In "Fish Physiology Vol IXB" Ed by WS Hoar, DJ Randall, EM Donaldson, Academic Press, New York, pp 117–170
- Goetz FW, Duman P, Ranjan M, Herman CA (1989) Prostaglandin F and E synthesis by specific tissue components of the brook trout (*Salvelinus fontinalis*) ovary. *J Exp Zool* 250: 196–205
- Goetz FW, Theofan G (1979) *In vitro* stimulation of germinal vesicle breakdown and ovulation of yellow perch (*Perca flavescens*) oocytes: effects of 17 $\alpha$ -hydroxy-20 $\beta$ -dihydroprogesterone and prostaglandins. *Gen Comp Endocrinol* 37: 273–285
- Greenhalgh EA (1990) Luteal steroidogenesis and regression in the rat: progesterone secretion and lipid peroxidation induced in luteal cells by human chorionic gonadotropin, phospholipase A<sub>2</sub> and prostaglandin F<sub>2 $\alpha$</sub> . *J Endocrinol* 125: 397–402
- Guillette LJ, Jr, Bjordal KA, Bolten AB, Gross TS, Palmer BD, Witherington BE, Matter JM (1991) Plasma estradiol-17 $\beta$ , progesterone, prostaglandin F, and prostaglandin E<sub>2</sub> concentrations during natural oviposition in the loggerhead turtle (*Caretta caretta*). *Gen Comp Endocrinol* 82: 121–130
- Guillette LJ, Jr, Cree A, Gross TS (1990) Endocrinology of oviposition in the tuatara (*Sphenodon punctatus*): I. Plasma steroids and prostaglandins during natural nesting. *Biol Reprod* 43: 285–289
- Guillette LJ, Jr, Lavia LA, Walker NJ, Roberts DK (1984) Luteolysis induced by prostaglandin F<sub>2 $\alpha$</sub>  in the lizard, *Anolis carolinensis*. *Gen Comp Endocrinol* 56: 271–277
- Ho SM (1987) Endocrinology of vitellogenesis. In "Hormones and Reproduction in Fishes, Amphibians, and Reptiles" Ed by DO Norris, RE Jones, Plenum Press, New York, pp 145–170
- Jones RE (1987) Ovulation: insights about the mechanisms based on a comparative approach. In "Hormones and Reproduction in Fishes, Amphibians, and Reptiles" Ed by DO Norris, RE Jones, Plenum Press, New York, pp 203–240
- Mahmoud IY, Cyrus RV, McAsey ME, Cady D, Woller MJ (1988) The role of arginine vasotocin and

- prostaglandin  $F_{2\alpha}$  on oviposition and luteolysis in the common snapping turtle *Chelydra serpentina* Gen Comp Endocrinol 69: 56-64
- 25 Ogata H, Nomura T, Hata M (1979) Prostaglandin  $PGF_{2\alpha}$  changes induced by ovulatory stimuli in the pond loach, *Misgurnus anguillicaudatus*. Bull Jpn Soc Sci Fish 45: 929-931
- 26 Schuetz AW (1967) Effect of steroids on germinal vesicle of oocytes of the frog (*Rana pipiens*) *in vitro*. Proc Soc Ex Biol Me 124: 1307-1310
- 27 Schuetz AW, Lessman AL (1981) Role of the surface epithelium of the ovarian follicle in the ovulation of frog (*Rana pipiens*) oocytes. J Cell Biol 91: 187a
- 28 Scossiroli RE, Palenzona DL (1979) Manuale di biometria. Zanichelli Editore, Bologna
- 29 Sedrani SH, El-Banna AA (1987) Effect of indomethacin on plasma level of vitamin D metabolites, oestradiol and progesterone in rabbit during early pregnancy. Comp Biochem Physiol 87: 635-639
- 30 Shimada K, Olson DM, Etches RJ (1984) Follicular and uterine prostaglandin levels in relation to uterine contraction and the first ovulation of a sequence in the hen. Biol Reprod 31: 76-82
- 31 Sokal RR, Rohlf FJ (1981) Biometry. Freeman, San Francisco, 2nd ed
- 32 Thibier-Fouchet C, Mulner O, Ozon R (1976) Progesterone biosynthesis and metabolism by ovarian follicles and isolated oocytes of *Xenopus laevis*. Biol Reprod 14: 317-326